



ELSEVIER

6 December 1999

PHYSICS LETTERS A

Physics Letters A 263 (1999) 341–346

www.elsevier.nl/locate/physleta

# Renormalization group analysis of the small-world network model

M.E.J. Newman<sup>\*</sup>, D.J. Watts

*Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA*

Received 16 August 1999; received in revised form 30 September 1999; accepted 21 October 1999

Communicated by A.R. Bishop

---

## Abstract

We study the small-world network model, which mimics the transition between regular-lattice and random-lattice behavior in social networks of increasing size. We contend that the model displays a critical point with a divergent characteristic length as the degree of randomness tends to zero. We propose a real-space renormalization group transformation for the model and demonstrate that the transformation is exact in the limit of large system size. We use this result to calculate the exact value of the single critical exponent for the system, and to derive the scaling form for the average number of ‘degrees of separation’ between two nodes on the network as a function of the three independent variables. We confirm our results by extensive numerical simulation. © 1999 Published by Elsevier Science B.V. All rights reserved.

*PACS:* 87.23.Ge; 05.10.Cc; 05.70.Jk; 64.60.Fr

*Keywords:* Small-world model; Renormalization group; Scaling; Phase transition

---

Unlike the lattices with which physics commonly deals, many real-world networks of interactions appear simultaneously to possess properties both of random graphs and of regular lattices. Watts and Strogatz [1] have recently proposed a network model that interpolates between these two extremes by taking a regular lattice and randomly ‘rewiring’ some of its edges. The resulting graphs are characterized by a high degree of local clustering (like regular lattices), but also possess short vertex–vertex distances similar to those found in random graphs even for quite

small densities of rewired edges. These ‘small-world’ networks, named after the small-world phenomenon of sociology [2], provide a model for the topology of a wide variety of systems, such as the Internet [3], power grids, patterns of neuron connectivity, and even networks of movie actors [4]. The structural properties of small-world networks have also been found to have a dramatic influence on dynamical systems [1], and may have significant consequences for many real-world applications, including coupled oscillators [5], neural networks, biological evolution [6], diffusion processes [7], and information propagation.

Intriguing though they are, the results of Watts and Strogatz are based primarily on computer simu-

---

<sup>\*</sup> Corresponding author. E-mail: mark@santafe.edu

lations, and our understanding of the small-world model stands to benefit from a more rigorous analysis. The techniques of statistical physics are especially appropriate to this task, particularly techniques drawn from the study of critical phenomena. In this paper, we argue that the small-world model has a critical point in the limit where the density of shortcuts tends to zero, and we investigate the model's behavior in the vicinity of this point using a renormalization group (RG) method, calculating the scaling forms and the single critical exponent describing the critical region.

Previous studies have concentrated on the one-dimensional version of the small-world model, and we will start with this version too, although we will later generalize our results to higher dimensions. In one dimension the model is defined on a lattice with  $L$  sites and periodic boundary conditions (the lattice is a ring). Initially each site is connected to all of its neighbors up to some fixed range  $k$  to make a network with coordination number  $z = 2k - 1$ . Randomness is then introduced by independently rewiring each of the  $kL$  connections with probability  $p$ . 'Rewiring' in this context means moving one end of the connection to a new, randomly chosen site. The behavior of the network thus depends on three independent parameters:  $L$ ,  $k$  and  $p$ . In this paper we will study a slight variation on the model in which shortcuts are added between randomly chosen pairs of sites, but no connections are removed from the regular lattice. For large  $L$ , this makes no difference to the mean separation between vertices of the network up to order  $p^{2k-1}$ . At order  $p^{2k}$  and higher it does make a difference, since the original small-world model is poorly defined in this case – there is a finite probability of a part of the lattice becoming disconnected from the rest and therefore making an infinite contribution to the average distance between vertices, and this makes the distance averaged over all networks also infinite. Our variation does not suffer from this problem and this makes the analysis significantly simpler. In Fig. 1 we show some examples of small-world networks.




Fig. 1. The RG transformations used in the calculations described in the text: (a) the transformation used for the  $k = 1$  system; (b) the transformation used for  $k > 1$ , illustrated in this case for  $k = 3$ . The shading of the sites indicates how they are grouped under the transformations.

We consider the behavior of the model for low density  $p$  of shortcuts. The fundamental observable quantity that we measure is the shortest distance between a pair of vertices on the network, averaged both over all pairs on the network and over all possible realizations of the randomness. This quantity, which we denote  $\ell$ , has two regimes of behavior. For systems small enough that there is much less than one shortcut on the lattice on average,  $\ell$  is dominated by the connections of the regular lattice and can be expected to increase linearly with system size  $L$ . As the lattice becomes larger with  $p$  held fixed, the average number of shortcuts will eventually become greater than one and  $\ell$  will start to scale as  $\log L$ . The transition between these two regimes takes place at some intermediate system size  $L = \xi$ , and from the arguments above we would expect  $\xi$  to take a value such that the number of shortcuts  $pk\xi \approx 1$ . In other words we expect  $\xi$  to diverge in the limit of small  $p$  as  $\xi \sim p^{-1}$ . The quantity  $\xi$  plays a role similar to the correlation length in an interacting system in conventional statistical physics, and its divergence leaves the small-world model with no characteristic length scale other than the fundamental lattice spacing. Thus the model possesses a phase transition at  $p = 0$ , and, as we will see, this gives rise to specific finite-size scaling

<sup>1</sup> In Ref. [1]  $k$  is defined to be equal to the coordination number  $z$ . Here we use  $k = \frac{1}{2}z$  to avoid unnecessary factors of 2 in our equations.

behavior in the region close to the transition <sup>2</sup>. Note that the phase transition at  $p = 0$  is a one-sided one, since  $p$  can never take a value less than zero. In this respect the transition is similar to transitions seen in other one-dimensional systems such as 1D bond or site percolation, or the 1D Ising model.

Barthélemy and Amaral [9] have suggested that the arguments above, although correct in outline, are not correct in detail. They contend that the length-scale  $\xi$  diverges as

$$\xi \sim p^{-\tau}, \quad (1)$$

with  $\tau$  different from the value of 1 given by the scaling argument. On the basis of numerical results, they conjecture that  $\tau = \frac{2}{3}$ . Barrat [10], on the other hand, has given a simple physical argument which directly contradicts this, indicating that  $\tau$  should be greater than or equal to 1. Amongst other things, we demonstrate in this paper that in fact  $\tau$  is exactly 1 for all values of  $k$  <sup>3</sup>.

Let us first consider the small-world model for the simplest case  $k = 1$ . As discussed above, the average distance  $\ell$  scales linearly with  $L$  for  $L \ll \xi$  and logarithmically for  $L \gg \xi$ . If  $\xi$  is much larger than one (i.e., we are close to the critical point), this implies that  $\ell$  should obey a finite-size scaling law of the form

$$\ell = Lf(L/\xi), \quad (2)$$

where  $f(x)$  is a universal scaling function with the limiting forms

$$f(x) \sim \begin{cases} \text{constant} & \text{for } x \ll 1 \\ (\log x)/x & \text{for } x \gg 1. \end{cases} \quad (3)$$

In fact, it is easy to show that the limiting value of  $f(x)$  as  $x \rightarrow 0$  is  $\frac{1}{4}$ . A scaling law similar though not identical to this has been proposed previously by

Barthélemy and Amaral [9] for the small-world model, although curiously they suggested that scaling of this type was evidence for the *absence* of a phase transition in the model, whereas we regard it as the appropriate form for  $\ell$  in the presence of one <sup>4</sup>.

We now assume that, in the critical region,  $\xi$  takes the form (1), and that we do not know the value of the exponent  $\tau$ . Then we can rewrite Eq. (3) in the form

$$\ell = Lf(p^\tau L), \quad (4)$$

where we have absorbed a multiplicative constant into the argument of  $f(x)$ , but otherwise it is the same scaling function as before, with the same limits, Eq. (3).

Now consider the real-space RG transformation on the  $k = 1$  small-world model in which we block sites in adjacent pairs to create a one-dimensional lattice of a half as many sites. (We assume that the lattice size  $L$  is even. In fact the transformation works fine if we block in groups of any size which divides  $L$ .) Two vertices are connected on the renormalized lattice if either of the original vertices in one was connected to either of the original vertices in the other. This includes shortcut connections. The transformation is illustrated in Fig. 1a for a lattice of size  $L = 24$ .

The number of shortcuts on the lattice is conserved under the transformation, so the fundamental parameters  $L$  and  $p$  renormalize according to

$$L' = \frac{1}{2}L, \quad p' = 2p. \quad (5)$$

The transformation generates all possible configurations of shortcuts on the renormalized lattice with the correct probability, as we can easily see since the probability of finding a shortcut between any two sites  $i$  and  $j$  is uniform, independent of  $i$  and  $j$  both before and after renormalization.

<sup>2</sup> Following the distribution of an early version of this paper, there has been some discussion in the literature concerning the exact nature of this transition. Although it seems widely accepted that the transition exists, its identification as first-order or continuous remains an open question. See [8].

<sup>3</sup> After seeing this paper in preprint form, as well as the preprint by Barrat, Ref. [10], Barthélemy and Amaral have conceded that indeed  $\tau = 1$ , and agree that the error arose from looking at systems too small to show the true scaling behavior (see Ref. citenote4).

<sup>4</sup> Ref. [9] is a little confusing in this respect. It appears the authors may have been referring to the possibility that the system shows a phase transition as the size  $L$  of the system is varied. This however would not be a sensible suggestion, since it is well-known that systems of finite size do not show sharp phase transitions. The only sensible scenario is a phase transition with varying shortcut probability  $p$ , which the model does indeed seem to show.

In almost all cases, the geometry of the shortest path between any two vertices is unchanged by our transformation, and it is straightforward to show that the number of vertex pairs for which the geometry does change is negligible for large  $L$  and small  $p$ . The length of a particular path is, on average, halved along those portions which run around the perimeter of the ring, and remains the same along the shortcuts. For large  $L$  and small  $p$ , the portion of the length along the shortcuts tends to zero and so can be neglected. Thus

$$\ell' = \frac{1}{2}\ell \quad (6)$$

in this limit.

Eqs. (5) and (6) constitute the RG equations for this system and are exact for  $L \gg 1$  and  $p \ll 1$ . Substituting into Eq. (4) we then find that

$$\tau = \frac{\log(L/L)}{\log(p'/p)} = 1. \quad (7)$$

Now we turn to the case of  $k > 1$ . To treat this case we define a slightly different RG transformation: we group adjacent sites in groups of  $k$ , with connections assigned using the same rule as before. The transformation is illustrated in Fig. 1b for a lattice of size  $L = 24$  with  $k = 3$ . Again the number of shortcuts in the network is preserved under the transformation, which gives the following renormalization equations for the parameters:

$$L' = L/k, \quad p' = k^2 p, \quad k' = 1, \quad \ell' = \ell. \quad (8)$$

Note that, in the limit of large  $L$  and small  $p$ , the mean distance  $\ell$  is not affected at all; the number of vertices along the path joining two distant sites is reduced by a factor  $k$ , but the number of vertices that can be traversed in one step is reduced by the same factor, and the two cancel out. For the same reasons as before, this transformation is exact in the limit of large  $L$  and small  $p$ .

We can use this second transformation to turn any network with  $k > 1$  into a corresponding network with  $k = 1$ , which we can then treat using the arguments given before. Thus, we conclude, the exponent

$\tau = 1$  for all values of  $k$  and, substituting from Eq. (8) into Eq. (4), the general small-world network must obey the scaling form

$$\ell = \frac{L}{k} f(pkL). \quad (9)$$

This form should be correct for  $L \gg 1$  and  $p' \ll 1$ , which implies that  $L/k \gg 1$  and  $k^2 p \ll 1$ . The first of these conditions is trivial – it merely ensures that we can neglect inaccuracies of  $\pm k$  in our estimate of  $\ell$  arising because positions on the lattice are rounded off to the nearest multiple of  $k$  by the RG transformation. The second condition is interesting however; it is necessary to ensure that the average distance traveled along shortcuts in the network is small compared to the distance traveled around the perimeter of the ring. This condition tells us when we are moving out of the scaling regime close to the transition, which is governed by (9), into the regime of the true random network, for which (9) is violated and  $\ell$  is known to scale as  $\log L / \log k$  [12]. It implies that we need to work with values of  $p$  which decrease as  $k^{-2}$  with increasing  $k$  if we wish to see clean scaling behavior, or conversely, that true random-network behavior should be visible in networks with values of  $p \simeq k^{-2}$  or greater.

We have tested our predictions by extensive numerical simulation of the small-world model. We have calculated exhaustively the minimum distance between all pairs of points on a variety of networks and averaged the results to find  $\ell$ . We have done this for  $k = 1$  (coordination number  $z = 2$ ) for systems of size  $L$  equal to a power of two from 128 up to 8192 and  $p = 1 \times 10^{-4}$  up to  $3 \times 10^{-2}$ , and for  $k = 5$  ( $z = 10$ ) with  $L = 512 \dots 32768$  and  $p = 1 \times 10^{-6} \dots 3 \times 10^{-4}$ . Each calculation was averaged over 1000 realizations of the randomness. In Fig. 2 we show our results plotted as the values of  $\ell k / L$  against  $pkL$ . Eq. (9) predicts that when plotted in this way the results should collapse onto a single curve and, as the figure shows, they do indeed do this to a reasonable approximation.

As mentioned above, Barthélemy and Amaral [9] also performed numerical simulations of the small-world model and extracted a value of  $\tau = \frac{2}{3}$  for the critical exponent. In the inset of Fig. 2 we show our simulation results for  $k = 1$  plotted according to Eq.




Fig. 2. Collapse of numerical data for  $\xi$  according to Eq. (9) as described in the text. Error bars are in all cases smaller than the data points. Note that the horizontal axis is logarithmic. Inset: the collapse is noticeably poorer for  $\tau = \frac{2}{3}$ .

(4) using this value for  $\tau$ . As the figure shows, the data collapse is significantly poorer in this case than for  $\tau = 1$ . It is interesting to ask then how Barthélemy and Amaral arrived at their result. It seems likely that the problem arises from looking at systems that are too small to show the true scaling behavior. In our calculations, we find good scaling for  $L/k \geq 60$ . Barthélemy and Amaral examined networks with  $k = 5, 10$  and  $15$  ( $z = 10, 20, 30$ ) so we should expect to find good scaling behavior for values of  $L$  larger than about 600. However, the systems studied by Barthélemy and Amaral ranged in size from about  $L = 50$  to about 500 in most cases, and in no case exceeded  $L = 1000$ . Their calculations therefore had either no overlap with the scaling regime, or only a small overlap, and so we would not expect to find behavior typical of the true value of  $\tau$  in their results (see footnote 3 and [11]).

It is possible to generalize the calculations presented here to small-world networks built on lattices of dimension  $d$  greater than one. For simplicity we consider first the case  $k = 1$ . If we construct a square or (hyper)cubic lattice in  $d$  dimensions with linear dimension  $L$ , connections between nearest-neighbor vertices, and shortcuts added with a rewiring probability of  $p$ , then as before the average vertex–vertex distance scales linearly with  $L$  for small  $L$ , logarithmically for large  $L$ , and the length-scale  $\xi$  of the

transition diverges according to Eq. (1) for small  $p$ . Thus the scaling form (4) applies for general  $d$  also. The appropriate generalization of our RG transformation involves grouping sites in square or cubic blocks of side 2, and the quantities  $L, p$  and  $\xi$  then renormalize according to

$$L' = \frac{1}{2}L, \quad p' = 2^d p, \quad \xi' = \frac{1}{2}\xi. \tag{10}$$

Thus

$$\tau = \frac{\log(L/L')}{\log(p'/p)} = \frac{1}{d}. \tag{11}$$

As an example, we show in Fig. 3 numerical results for the  $d = 2$  case, for  $L$  equal to a power of two from 64 up to 1024 (i.e., a little over a million vertices for the largest networks simulated) and six different values of  $p$  for each system size from  $p = 3 \times 10^{-6}$  up to  $1 \times 10^{-3}$ . The results are plotted according to Eq. (4) with  $\tau = \frac{1}{2}$  and, as the figure shows, they again collapse nicely onto a single curve.

A number of generalizations are possible for  $k > 1$ . Perhaps the simplest is to add connections along the principal axes of the lattice between all vertices whose separation is  $k$  or less. This produces a graph with average coordination number  $z = 2dk$ . By blocking vertices in square or cubic blocks of edge  $k$ , we can then transform this system into one with  $k = 1$ . The appropriate generalization of the RG Eq.




Fig. 3. Collapse of numerical data for networks based on the square lattice in two dimensions, as described in the text. Error bars are in all cases smaller than the data points.

(8) is then

$$L' = L/k, \quad p' = k^{d+1}p, \quad k' = 1, \quad \ell' = \ell, \quad (12)$$

which gives  $\tau = 1/d$  for all  $k$  and a scaling form of

$$\ell = \frac{L}{k} f((pk)^{1/d} L). \quad (13)$$

Alternatively, we could redefine our scaling function  $f(x)$  so that  $\ell k/L$  is given as a function of  $pkL^d$ . Writing it in this form makes it clear that the number of vertices in the network at the transition from large- to small-world behavior diverges as  $p^{-1}$  in any number of dimensions.

Another possible generalization to  $k > 1$  is to add connections between all sites within square or cubic regions of side  $2k$ . This gives a different dependence on  $k$  in the scaling form, but  $\tau$  still equal to  $1/d$ .

To conclude, we have studied the small-world network model of Watts and Strogatz using an asymptotically exact real-space renormalization group method. We find that in all dimensions  $d$  the model has a critical point in the limit where the density of shortcuts  $p$  tends to zero and that the characteristic length  $\xi$  diverges according to  $\xi \sim p^{-\tau}$  with  $\tau = 1/d$  for all values of the connection range  $k$ . We have also deduced the general finite-size scaling law which describes the variation of the mean vertex–vertex separation as a function of  $p$ ,  $k$  and the system size  $L$ . We have performed extensive numerical calculations which confirm our analytic results.

## Acknowledgements

This work was supported in part by the Santa Fe Institute and by funding from the NSF (grant number PHY-9600400), the DOE (grant number DE-FG03-94ER61951), and DARPA (grant number ONR N00014-95-1-0975).

## References

- [1] D.J. Watts, S.H. Strogatz, *Nature* 393 (1998) 440.
- [2] S. Milgram, *Psychol. Today* 2 (1967) 60.
- [3] L. Adamic, Xerox PARC working paper <http://www.parc.xerox.com/istl/groups/iea/www/SmallWorld.html>, 1999.
- [4] D.J. Watts, *Small Worlds: The Dynamics of Networks Between Order and Randomness*, Princeton University Press, Princeton, 1999.
- [5] Y. Kuramoto, *Chemical Oscillations, Waves and Turbulence*, Springer, Berlin, 1984.
- [6] R.V. Kulkarni, E. Almaas, D. Stroud, *cond-mat/9905066*.
- [7] R. Monasson, submitted to *Eur. Phys. J. B*. Also *cond-mat/9903347*.
- [8] M. Argollo de Menezes, C.F. Moukarzel, T.J.P. Penna, First-order transition in small-world networks, submitted to *Phys. Rev. Lett*. Also *cond-mat/9903426*.
- [9] M. Barthélemy, L.A.N. Amaral, *Phys. Rev. Lett.* 82 (1999) 3180.
- [10] A. Barrat, submitted to *Phys. Rev. Lett*. Also *cond-mat/9903323*.
- [11] M. Barthélemy, L.A.N. Amaral, *Phys. Rev. Lett.* 82 (1999) 5180.
- [12] B. Bollobás, *Random Graphs*, Academic Press, New York, 1985.